

Rodotet®



Medicated Premix

Active substance

Rodotet® premix contains two active substances: tiamulin hydrogen fumarate (THF) and chlortetracycline (CTC).

Indications

- Prevention and treatment of *Mycoplasma hyopneumoniae*, *Actinobacillus pleuropneumoniae*, *Brachyspira hyodysenteriae*, PRDC, *Lawsonia intracellularis* and *Spirochetosis*.

Target Species

Pigs

Dosage and Mode of administration

PRACTICAL ADMINISTRATION		
TYPE OF APPLICATION	RODOTET®	AS ACTIVE INGREDIENTS
PREVENTION	1.0 – 1.5 kg/ ton feed During 10 – 14 days	THF: 33-50 ppm CTC: 100-150 ppm
TREATMENT	3.0 – 4.5 kg/ ton feed During 7 days	THF: 100-150 ppm CTC: 300-450 ppm



Withdrawal Period

Meat and offal

- when used for treatment, for meat – 10 days after the last administration for preparation;
- when used for prevention, for meat – 1 day after the last administration for preparation.

Shelf Life

- Shelf-life of the veterinary medicinal product as packaged for sale: 2 years.
- Shelf-life after dilution in feed according to directions: 3 months.

Package

Sachets of 30 g, 150 g or 1 kg;

Drums of 5 kg, 10 kg or 25 kg

Storage conditions: Rodotet® Premix has a pale yellow to yellow color and is meant for oral use only. The product must be stored in its original packaging. The original packaging should be tightly closed.

Exposure to direct sunlight shall be avoided.

