



Composition

Active substance: **800 mg amoxicillin trihydrate per g** (equivalent to 697 mg amoxicillin per g)

Pharmaceutical form

Powder for use in drinking water. White to slightly yellow powder.

Target species

- Pigs
- Chickens (broilers, pullets, breeders)
- Turkeys (broilers, breeders)
- Ducks (broilers, breeders)

Indications

Pigs

- Treatment of *Pasteurella multocida* susceptible to amoxicillin.

Chickens, turkeys and ducks

- Treatment of infections caused by bacteria susceptible to amoxicillin.

Practical dosing and administration

In drinking water or in liquid feed (pigs)

Pigs

- 20 mg amoxicillin trihydrate/kg bw/day, equivalent to 17.4 mg of amoxicillin/kg bw/day (corresponding to 25 mg product/kg bw/day)
- Up to 5 days

Chickens

- 15 mg amoxicillin trihydrate per kg bw/day, equivalent to 13.1 mg of amoxicillin/kg bw/day (corresponding to 18.8 mg product/kg bw/day)
- For 3 days or in severe cases for 5 days

Turkeys

- 15-20 mg amoxicillin trihydrate/kg bw/day, equivalent to 13.1-17.4 mg of amoxicillin/kg bw/day (corresponding to 18.8-25 mg product/kg bw/day)
- For 3 days or in severe cases for 5 days

Ducks

- 20 mg amoxicillin trihydrate/kg bw/day, equivalent to 17.4 mg of amoxicillin/kg bw/day (corresponding to 25 mg product/kg bw/day)
- For 3 consecutive days

Drinking water

The complete dissolution of the product should be ensured, by mixing gently the solution up to complete dissolution. The maximum solubility of the product in water is 8 g/L at 20°C and 3 g/L at 5°C. Any medicated water which is not consumed within 24 hours should be discarded and the medicated drinking water replenished.

Liquid feed

Medicated feed should be freshly prepared on at least 2 occasions per day over the treatment period. In order to ensure that any loss of amoxicillin activity is minimized, the quantity of medicated liquid feed prepared should not exceed the amount of feed which will be consumed within 2 hours.

Overdose

No problems with overdosage have been reported. Treatment should be symptomatic and no specific antidote is available.

Withdrawal period(s)

Meat and offal

- Pigs: 2 days
- Chickens: 1 days
- Turkeys: 5 days
- Ducks: 9 days

Not authorised for use in laying birds producing eggs for human consumption. Do not use within 4 weeks of the start of the laying period.

Shelf life

- As packaged for sale: 2 years
- After first opening the immediate packaging: 6 months
- After dissolution in drinking water according to directions: **24 hours**
- After incorporation in liquid feed according to directions: 2 hours

Storage

- Do not refrigerate or freeze.
- Store in the original container in order to protect from light.
- Keep the container tightly closed.
- Store in a dry place.

Packaging

- Zipped 500 g and 1 kg bag made of low density polyethylene/ aluminium/polyethylene terephthalate.
- 100 g jar made of high density polyethylene closed with a seal made of low density polyethylene/ polyethylene terephthalate/ aluminium and a screw cap made of polypropylene.
- Thermo-sealed 100 g bag made of low density polyethylene/ aluminium/polyethylene terephthalate.

Not all pack sizes may be marketed.



Contraindications

Do not use in horses, rabbits, guinea pigs, hamsters, gerbils or any other small herbivore given that amoxicillin, as for all aminopenicillins, has a deleterious effect on caecal bacteria. Do not use in adult ruminants. Do not use in animals with known hypersensitivity to penicillins or other β -lactam antibiotics or to any of the excipients. Do not administer to animals with renal disease including anuria or oliguria. Do not use in the presence of β -lactamase-producing bacteria.

- Full SPC can be obtained on request
- Indications listed above are not necessarily authorized in all countries. Please consult the local label for exact indications and posology.
- POM-V
- VM 30282/4046



Huvamox®