

## Pharmasin® 100 WSG

### Water soluble granules



### Active substance

- **1.1 g of granules contains 1 g of tylosin (corresponding to 1.1 g of tylosin tartrate)**

### Indications

#### Calves

- Treatment and prevention of pneumonia caused by *Mycoplasma* spp when the disease has been established in the herd.

#### Pigs

- Treatment and prevention of enzootic pneumonia caused by *Mycoplasma hyopneumoniae* and *Mycoplasma hyorhinis* when the disease has been established in the herd.
- Treatment and prevention of *Porcine Intestinal Adenomatosis* (Ileitis) associated with *Lawsonia intracellularis* when the disease has been established in the herd.

#### Chickens

- Treatment and prevention of chronic respiratory diseases (CRD) caused by *Mycoplasma gallisepticum* and *Mycoplasma synoviae* when the disease has been established in the flock.
- Treatment and prevention of necrotic enteritis caused by *Clostridium perfringens* when the disease has been established in the flock.

#### Turkeys

- Treatment and prevention of infectious sinusitis caused by *Mycoplasma gallisepticum* when the disease has been established in the flock.

### Target Species

Calves, Pigs, Chickens and Turkeys

### Dosage and Mode of administration

Oral administration through the drinking water. In calves the product can also be administered through milk or milk replacer.

#### Calves

- 10-20 mg tylosin per kg BW (corresponding to 11-22 mg of the veterinary medicinal product per kg BW ), twice daily (corresponding to a daily dose of 20-40 mg tylosin per kg BW), for 7-14 days.

#### Pigs

- For the treatment of enzootic pneumonia: 20 mg tylosin per kg BW per day (corresponding to 22 mg of the veterinary medicinal product per kg BW ) for 10 days.
- For the treatment of ileitis or PIA: 5-10 mg tylosin per kg BW per day (corresponding to 5.5-11 mg of the veterinary medicinal product per kg BW ) for 7 days.

#### Chickens

- For the treatment of chronic respiratory disease: 75-100 mg tylosin per kg BW per day (corresponding to 82.5-110 mg of the veterinary medicinal product per kg BW ) for 3-5 days.
- For the treatment of necrotic enteritis: 20 mg tylosin per kg BW per day (corresponding to 22 mg of the veterinary medicinal product) for 3 days.

#### Turkeys

- 75-100 mg tylosin per kg BW per day corresponding to 82.5-110 mg of the veterinary medicinal product per kg BW ) for 3-5 days.

### Withdrawal Period

- |                              |         |
|------------------------------|---------|
| • Calves (meat and offal):   | 12 days |
| • Pigs (meat and offal):     | 1 day   |
| • Turkeys (meat and offal):  | 2 days  |
| • Turkey (eggs):             | 0 days  |
| • Chickens (meat and offal): | 1 day   |
| • Chicken (eggs):            | 0 days  |

### Shelf Life

- Shelf-life of the veterinary medicinal product as packaged for sale: 3 years
- Shelf-life after dilution or reconstitution according to directions:
  - medicated water: 24 hours
  - medicated milk or milk replacer: 24 hours
- Shelf-life after first opening the immediate packaging: 3 months

### Packaging

- 1.1 kg resealable block bottom zipped sachet made of polyethylene/aluminium/polyethylene terephthalate laminate
- 110 g high density polyethylene pot with polypropylene cap

